Appendix A

Claim Amendments

1. (Currently amended) A compound of formula 1,

$$R1$$
 H $R4$ $R4$ $R3$ $R4$ $R4$ $R4$ $R5$ $R5$ $R5$ $R5$

in which

R1 is a mono- or bicyclic aromatic radical substituted by R11, R12, R13 and R14, wherein R1 is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furanyl (furyl), benzofuranyl (benzofuryl), thiophenyl (thienyl), benzothiophenyl (benzothienyl), thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, quinolinyl and isoquinolinyl, where

- is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-R11 4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, aminocarbonyl, di-1-4Ccarboxyl, monooralkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkoxycarbonyl-1-4C-alkyl, 1-4C-alkyl, hydroxyl, aryl, aryl-1-4C-alkyl, halogen, aryloxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, di-1-4C-alkylamino, amino, monoor1-4Calkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl, together with R12 methylenedioxy or ethylenedioxy,
- R12 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydroxyl, or together with R11 methylenedioxy or ethylenedioxy,
- R13 is hydrogen, 1-4C-alkyl or halogen and R14 is hydrogen, 1-4C-alkyl or halogen, where
- aryl is phenyl or substituted phenyl having one, two or three substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, carboxyl,

1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxyl, and cyano and mixtures thereof,

is a mono- or bicyclic aromatic radical substituted R2 by R21, R22, R23 and R24, wherein R2 is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furanyl (furyl), benzofuranyl (benzofuryl), thiophenyl (thienyl), benzothiophenyl (benzothienyl), thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, quinolinyl and isoquinolinyl, where is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-R21 4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, mono- or di-1-4Calkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, aryl, halogen, hydroxyl, aryl-1-4C-alkyl, aryloxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, di-1-4C-alkylamino, 1-4Camino, monoor alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl, methylenedioxy together with R22 or

ethylenedioxy,

- R22 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydroxyl, or together with R21 methylenedioxy or ethylenedioxy,
- R23 is hydrogen, 1-4C-alkyl or halogen and R24 is hydrogen, 1-4C-alkyl or halogen, where
- aryl is phenyl or substituted phenyl having one, two or three substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, carboxyl, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxyl, and cyano and mixtures thereof,
- is a mono- or bicyclic aromatic radical substituted by R31, R32, R33 and R34, wherein R3 is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furanyl (furyl), benzofuranyl (benzofuryl), thiophenyl (thienyl), benzothiophenyl (benzothienyl), thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, quinolinyl and isoquinolinyl, where

- is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-R31 1-4C-alkylcarbonyl, alkoxy, 2-4C-alkenyloxy, aminocarbonyl, di-1-4Ccarboxyl, monooralkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkoxycarbonyl-1-4C-alkyl, 1-4C-alkyl, halogen, hydroxyl, aryl, aryl-1-4C-alkyl, aryloxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4Calkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl, ortogether with R32 methylenedioxy ethylenedioxy,
- R32 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydroxyl, or together with R31 methylenedioxy or ethylenedioxy,
- R33 is hydrogen, 1-4C-alkyl or halogen and R34 is hydrogen, 1-4C-alkyl or halogen, where
- aryl is phenyl or substituted phenyl having one, two or three substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, carboxyl,

1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxyl, and cyano and mixtures thereof,

R4 is hydrogen or methyl,

R5 is hydrogen or methyl,

Al is 1-3C-alkylene or ethyleneoxy (-CH₂-CH₂-O-) and

A2 is 1-3C-alkylene or ethyleneoxy (-CH₂-CH₂-O-),

or a hydrate, solvate, salt, hydrate of a salt or solvate

of a salt thereof.

 (Currently amended) A compound of formula 1 according to claim 1,

in which

is a mono- or bicyclic aromatic radical substituted by R11, R12, R13 and R14, wherein R1 is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furanyl (furyl), benzofuranyl (benzofuryl), thiophenyl (thienyl), benzothiophenyl (benzothienyl), thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, quinolinyl and isoquinolinyl, where

R11 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-

2-4C-alkenyloxy, 1-4C-alkylcarbonyl, alkoxy, carboxyl, aminocarbonyl, monoordi-1-4Calkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkoxycarbonyl-1-4C-alkyl, 1-4C-alkyl, hydroxyl, halogen, aryl, aryl-1-4C-alkyl, aryl-1-4C-alkoxy, trifluoromethyl, aryloxy, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4Calkoxycarbonylamino, 1-4C-alkoxy-1-4Calkoxycarbonylamino or sulfonyl,

- R12 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxy, alkoxycarbonyl, halogen, trifluoromethyl or hydroxyl,
- R13 is hydrogen, 1-4C-alkyl or halogen and R14 is hydrogen, 1-4C-alkyl or halogen, where
- aryl is phenyl or substituted phenyl having one, two or three substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, carboxyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxyl, and cyano and mixtures-thereof,

- is a mono- or bicyclic aromatic radical substituted by R2 R21, R22, R23 and R24, wherein R2 is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furanyl (furyl), benzofuranyl (benzofuryl), thiophenyl (thienyl), benzothiophenyl (benzothienyl), thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, quinolinyl and isoquinolinyl, where
 - is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-R21 alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, monodi-1-4Coralkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkoxycarbonyl-1-4C-alkyl, 1-4C-alkyl, halogen, hydroxyl, aryl, aryl-1-4C-alkyl, aryloxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, amino, di-1-4C-alkylamino, monoor alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino orsulfonyl,
 - R22 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydro-

xyl,

R23 is hydrogen, 1-4C-alkyl or halogen and R24 is hydrogen, 1-4C-alkyl or halogen, where

- aryl is phenyl or substituted phenyl having one, two or three substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, carboxyl, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxyl, and cyano and mixtures—thereof,
- is a mono- or bicyclic aromatic radical substituted R3 by R31, R32, R33 and R34, wherein R3 is selected from the group consisting of phenyl, naphthyl, imidazolyl, 1,2,3-triazolyl, pyrrolyl, pyrazolyl, indolyl, benzimidazolyl, furanyl (furyl), benzofuranyl (benzofuryl), thiophenyl (thienyl), benzothiophenyl (benzothienyl), thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, quinolinyl and isoquinolinyl, where is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-R31 alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, monoordi-1-4C-

alkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-

1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxyl, aryl, aryl-1-4C-alkyl, aryloxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl,

R32 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydroxyl,

R33 is hydrogen, 1-4C-alkyl or halogen and R34 is hydrogen, 1-4C-alkyl or halogen, where

aryl is phenyl or substituted phenyl having one, two or three substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, carboxyl, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxyl, and cyano and mixtures thereof,

R4 is hydrogen,

R5 is hydrogen,

A1 denotes 1-3C-alkylene and

A2 denotes 1-3C-alkylene,

or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

 (Currently amended) A compound of formula 1 according to claim 1,

in which

- R1 is an aromatic radical substituted by R11, R12, R13 and R14, wherein R1 is selected from the group consisting of phenyl, furanyl (furyl) and thiophenyl (thienyl), where
 - R11 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, mono- or di-1-4Calkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4Chalogen, hydroxyl, trifluoromethyl, alkyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4Calkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl, together with R12 methylenedioxy or ethylenedioxy,
 - R12 is hydrogen or halogen, or together with R11

methylenedioxy or ethylenedioxy,

R13 is hydrogen and

R14 is hydrogen,

R2 is an aromatic radical substituted by R21, R22, R23 and R24, wherein R2 is selected from the group consisting of pyridinyl and pyrimidinyl,

where

R21 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy or halogen,

R22 is hydrogen or halogen,

R23 is hydrogen and

R24 is hydrogen,

R3 is an aromatic radical substituted by R31, R32, R33 and R34, wherein R3 is selected from the group consisting of phenyl and pyridinyl,

where

R31 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxyl, trifluoromethyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylamino, 1-4C-alkylamino,

alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl,

R32 is hydrogen or halogen,

R33 is hydrogen and

R34 is hydrogen,

R4 is hydrogen or methyl,

R5 is hydrogen or methyl,

A1 is 1-3C-alkylene or ethyleneoxy (-CH2-CH2-O-) and

A2 is 1-3C-alkylene or ethyleneoxy (-CH₂-CH₂-O-),

or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

- 4. (Currently amended) A compound of formula 1 according to claim 1, in which
- R1 is an aromatic radical substituted by R11, R12, R13 and R14, wherein R1 is selected from the group consisting of phenyl, furanyl (furyl) and thiophenyl (thienyl),

where

R11 is hydrogen, 1-4C-alkoxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, halogen, hydroxyl or mono- or di-1-4C-alkylamino, or together with R12

methylenedioxy or ethylenedioxy,

R12 is hydrogen or halogen, or together with R11 methylenedioxy or ethylenedioxy,

R13 is hydrogen and

R14 is hydrogen,

R2 is an aromatic radical substituted by R21, R22, R23 and R24, wherein R2 is selected from the group consisting of pyridinyl and pyrimidinyl, where

R21 is hydrogen,

R22 is hydrogen,

R23 is hydrogen and

R24 is hydrogen,

R3 is an aromatic radical substituted by R31, R32, R33 and R34, wherein R3 is selected from the group consisting of phenyl and pyridinyl, where

R31 is hydrogen, 1-4C-alkoxy or halogen,

R32 is hydrogen,

R33 is hydrogen and

R34 is hydrogen,

R4 is hydrogen or methyl,

R5 is hydrogen or methyl,

Al denotes methylene, ethylene, ethylidene [-CH(CH3)-] or

ethyleneoxy $(-CH_2-CH_2-O-)$ and

- A2 denotes methylene, ethylene, ethylidene [-CH(CH₃)-] or ethyleneoxy (-CH₂-CH₂-O-),
- or a hydrate, solvate, salt, hydrate of a salt or solvate

 of a salt thereof.
- 5. (Currently amended) A compound of formula 1 according to claim 1,

in which

R1 is an aromatic radical substituted by R11, R12, R13 and R14, wherein R1 is selected from the group consisting of phenyl, furanyl (furyl) and thiophenyl (thienyl),

where

- R11 is hydrogen, 1-4C-alkoxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, halogen, hydroxyl or mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, or together with R12 methylenedioxy or ethylenedioxy,
- R12 is hydrogen or halogen, or together with R11 methylenedioxy or ethylenedioxy,

R13 is hydrogen and R14 is hydrogen,

R2 is an aromatic radical substituted by R21, R22, R23 and R24, wherein R2 is selected from the group consisting of pyridinyl and pyrimidinyl, where

R21 is hydrogen,

R22 is hydrogen,

R23 is hydrogen and

R24 is hydrogen,

R3 is an aromatic radical substituted by R31, R32, R33 and R34, wherein R3 is selected from the group consisting of phenyl and pyridinyl, where

R31 is hydrogen, 1-4C-alkoxy or halogen,

R32 is hydrogen,

R33 is hydrogen and

R34 is hydrogen,

R4 is hydrogen or methyl,

R5 is hydrogen or methyl,

Al denotes methylene, ethylene, ethylidene (-CH(CH $_3$)-] or ethyleneoxy (-CH $_2$ -CH $_2$ -O-) and

A2 denotes methylene, ethylene, ethylidene [-CH(CH₃)-] or ethyleneoxy (-CH₂-CH₂-O-),

or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

- 6. (Currently amended) A compound of formula 1 according to claim 1, in which
- R1 is furanyl (furyl), thiophenyl (thienyl) or phenyl substituted by R11 and R12, where
 - R11 is hydrogen, 1-4C-alkoxy, 1-4C-alkylcarbonyl, carboxyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, halogen, hydroxyl or mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, or together with R12 methylenedioxy or ethylenedioxy,
 - R12 is hydrogen or halogen, or together with R11 methylenedioxy or ethylenedioxy,

R2 is pyridinyl,

R3 is phenyl,

R4 is hydrogen,

R5 is hydrogen,

Al denotes methylene and

A2 denotes methylene,

or a hydrate, solvate, salt, hydrate of a salt or solvate

of a salt thereof.

- 7. (Currently amended) A compound of formula 1 according to claim 1, in which
- R1 is furanyl (furyl), thiophenyl (thienyl) or phenyl substituted by R11 and R12,

where

R11 is hydrogen, 1-4C-alkoxy, carboxyl, aminocarbonyl, halogen or di-1-4C-alkylamino and R12 is hydrogen,

R2 is pyridinyl,

R3 is phenyl,

R4 is hydrogen,

R5 is hydrogen,

Al denotes methylene and

A2 denotes methylene,

or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

8. (Currently amended) A compound of formula 1 according to claim 1, wherein R2 is 2-pyridinyl or 4-pyridinyl, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

- 9. (Currently amended) A compound of formula 1 according to claim 1, wherein R3 is phenyl, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.
- 10. (Currently amended) A compound of formula 1 according to claim 1, wherein R1 is furanyl (furyl), thiophenyl (thienyl) or phenyl substituted by R11 and R12, where R11 is hydrogen, 1-4C-alkoxy, carboxyl, aminocarbonyl, halogen or di-1-4C-alkylamino and R12 is hydrogen, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.
- 11. (Currently amended) A compound of formula 1 according to claim 1, wherein R1 is furanyl (furyl), thiophenyl (thienyl) or phenyl substituted by R11 and R12, where R11 is 1-4C-alkoxy, halogen, di-1-4C-alkylamino, aminocarbonyl, carboxyl, 1-4C-alkylcarbonylamino, 1-4C-alkylcarbonyl hydroxyl, together orwith methylenedioxy or ethylenedioxy and R12 is hydrogen, halogen together with R11 or methylenedioxy ethylenedioxy, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

- 12. (Currently amended) A compound of formula 1 according to claim 1, wherein R1 is 2-furanyl or 3-furanyl, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.
- 13. (Currently amended) A compound of formula 1 according to claim 1, wherein R1 is 2-thiophenyl or 3-thiophenyl, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.
- 14. (Currently amended) A compound of formula 1 according to claim 1, wherein R1 is selected from the group consisting phenyl, 4-methoxyphenyl, 4-chlorophenyl, of dimethylaminophenyl, 4-aminocarbonylphenyl, 4 carboxyphenyl, 3-chloro-4-fluorophenyl, 3 acetylaminophenyl, benzo[1,3]dioxol-5-yl, 3-hydroxyphenyl, 4-hydroxyphenyl, 4-acetylphenyl, 3-acetylphenyl, acetylaminophenyl, 4-dimethylaminocarbonyl-phenyl and 4aminocarbonylphenyl, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.
- 15. (Currently amended) A compound of formula 1 according

to claim 1, wherein Al denotes methylene and A2 denotes methylene, or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof.

- 16. (Currently amended) A pharmaceutical composition comprising a compound of the formula 1 as claimed in claim 1 and/or a pharmaceutically acceptable hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof together with a pharmaceutically acceptable excipient and/or carrier.
- 17. (Currently amended) A method of treating a patient afflicted with a disease or disorder, comprising the step of administering a therapeutically effective amount of a compound of the formula 1 as claimed in claim 1 and/or a pharmaceutically acceptable hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof to said patient afflicted with said disease or disorder, wherein the disease or disorder is selected from the group consisting of an acute and/or chronic airway disorder, an inflammatory or allergen induced airway disorder, bronchitis, obstructive bronchitis, spastic bronchitis, allergic bronchitis,

allergie asthma, bronchial asthma, emphysema, chronic obstructive pulmonary disease (COPD), a disorder which is based on an excessive release of T Cell derived cytokines, HIV infection, septic shock, adult respiratory distress syndrome, graft versus host reactions, acute or chronic rejection of organ or tissue allo or xenografts, generalized inflammations in the gastrointestinal area, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, faulty immunological reactions in the area of the upper airways and the adjacent regions, dermatoses of the proliferative, inflammatory or allergic type, psoriasis (vulgaris), toxic and allergic contact eczema, atopic eczema, seborrheic eczema, lichen simplex, sunburn, pruritus in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, follicular and wide area pyodermias, endogenous and exogenous acne, acne rosacea, other proliferative, inflammatory, allergic skin disorders, a disorder in connection with disturbances of brain metabolism, disorders of the central nervous system (CNS), cerebral senility, senile dementia, multiinfarct dementia, depression, arteriosclerotic dementia, cancer and diabetes insipidus.

- 18. (Currently amended) A method according to claim 17, in which a compound of claim 1 and/or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof, is used in combination with other therapeutic agents used in clinical practice for the treatment of the said disease or disorder.
- 19. (Currently amended) A process for preparing a compound of formula 1 as claimed in claim 1 or a hydrate, solvate, salt, hydrate of a salt or solvate of a salt thereof, which comprises reacting a compound boronic acid derivative R1-B(OH)₂ wherein R1 has the meaning specified in claim 1, with a compound pyrimidine derivate of formula (4)

in which Al, A2, R2, R3, R4 and R5 have a meaning specified

in claim 1, and optionally converting an obtained compound into a corresponding hydrate, solvate, salt, hydrate of a salt or solvate of a salt, or converting an obtained hydrate, solvate, salt, hydrate of a salt or solvate of a salt into a corresponding free compound.